

## **PRACTICAL MANUAL**

# **Pests of crops and stored grains and their management**

**(ENT- 311) 3(2+1)**

**For B.Sc. (Hons.) Agriculture V Semester**

**Dr. Dharm Raj Singh**

**Dr. Ram Singh**

**Dr. Akshay Kumar**



**2022**

**Department of Entomology**

**College of Agriculture**

**Chandra Shekhar Azad University of Agriculture and  
Technology, Kanpur**

**Syllabus ENT-321 3(2+1):** Identification of different types of damage. Identification and study of life cycle and seasonal history of various insect pests attacking crops and their produce: (a) Field Crops; (b) Vegetable Crops; (c) Fruit Crops; (d) Plantation, gardens, Narcotics, spices & condiments. Identification of insect pests and Mites associated with stored grain. Determination of insect infestation by different methods. Assessment of losses due to insects. Calculations on the doses of insecticides application technique. Fumigation of grain store / godown. Identification of rodents and rodent control operations in godowns. Identification of birds and bird control operations in godowns. Determination of moisture content of grain. Methods of grain sampling under storage condition. Visit to Indian Storage Management and Research Institute, Hapur and Quality Laboratory, Department of Food., Delhi. Visit to nearest FCI godowns.

**Name of Student** .....

**Id.No.** .....

**Batch** .....

**Session** .....

**Semester** .....

**Course Name :** .....

**Course No. :** .....

**Credit** .....

---

### CERTIFICATE

This is to certify that Shri./Km. ....ID  
No..... has completed the practical of  
course.....course No. .... as per  
the syllabus of B.Sc. (Hons.) Agriculture/ Horticulture/ Forestry ..... semester in the  
year.....in the respective lab/field of College.

Date:

Course Teacher

## CONTENTS

<b>Sl. No.</b>	<b>Name of the EXERCISE</b>	<b>Page No.</b>
1	To study damage done by phytophagous insect	
2	To identify insect pests attacking cereal crops	
3	To identify insect pests attacking pulse crops	
4	To identify insect pests attacking cash crops	
5	To identify insect pests attacking vegetable crops	
6	To identify insect pests attacking oilseed crops	
7	To identify insect pests attacking fruit crops	
8	To identify insect pests attacking plantation crops	
9	To identify insect pests attacking spices, condiments, and narcotics	
10	To identify insect pests attacking ornamental plants	
11	To identify insect pests and Mites associated with stored grain	
12	To determine insect infestation	
13	To assess losses due to insects	
14	To calculate doses of insecticides application technique	
15	To study fumigation for stored grains	
16	To study non-insect pests in field/godown	
17	To determine the moisture content of grain	
18	To study methods of grain sampling under storage conditions.	
19	To visit Indian Storage Management and Research Institute, Hapur and Quality Laboratory, Department of Food., Delhi. Visit nearest FCI godowns.	

**EXERCISE NO. 1**

**Objective: To study damage done by phytophagous insect**

**Exercise:** Observe and write the damaging symptoms of the given insect specimen.

**OBSERVATIONS TO BE RECORDED:**

**Stem borer:** .....  
.....  
.....  
.....

Eg. ....  
.....  
.....

**Shoot borer:** .....  
.....  
.....

Eg. ....

**Defoliator:** .....  
.....  
.....

Eg. ....

**Leaf miner:**.....  
.....  
.....

Eg. ....

**Leaf Webber:**.....  
.....  
.....

Eg. ....

**Leaf folder:**.....  
.....  
.....

Eg. ....

**Gall maker:** .....

.....

Eg. ....

**Pod/ capsule borers/ bollworm:**.....

.....

.....

Eg. ....

**Root feeder:**.....

.....

.....

Eg. ....

**Seed feeder (stored grain pests) .....**

.....

.....

Eg. ....

**Sap feeder**

a. From grain: .....

.....

.....

Eg. ....

b. From tender plant parts:

.....

.....

.....

Eg. ....

**EXERCISE NO. 2**

**Objective: To identify insect pests attacking cereal crops**

**Exercise:** Observe and enlist major pests attacking cereal crops. Draw a neat diagram of important pests of cereal crops.

Common name	Scientific name	Family & Order	Pest identification/ Bionomics	Damaging symptoms
<b>Rice</b>				
Rice stem borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Rice gall midge	..... .....	..... .....	..... ..... .....	..... ..... .....
Green leaf hopper	..... .....	..... .....	..... ..... .....	..... ..... .....
Brown plant hopper	..... .....	..... .....	..... ..... .....	..... ..... .....
White back plant hopper	..... .....	..... .....	..... ..... .....	..... ..... .....

Rice earhead bug	..... .....	..... .....	..... ..... .....	..... ..... .....
Rice leaf folder	..... .....	..... .....	..... ..... .....	..... ..... .....
Rice caseworm	..... .....	..... .....	..... ..... .....	..... ..... .....
Rice grasshopper	..... .....	..... .....	..... ..... .....	..... ..... .....
<b>Wheat and Barley</b>				
Wheat aphid	..... .....	..... .....	..... ..... .....	..... ..... .....
Climbing cutworm /armyworm	..... .....	..... .....	..... ..... .....	..... ..... .....
Ghujhia Weevil	..... .....	..... .....	..... ..... .....	..... ..... .....

			.....	.....
Termites	..... .....	..... .....	..... ..... .....	..... ..... .....
<b>Maize and Sorghum</b>				
Stem borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Fall armyworm	..... .....	..... .....	..... ..... .....	..... ..... .....
Shoot fly	..... .....	..... .....	..... ..... .....	..... ..... .....
Shoot bug	..... .....	..... .....	..... ..... .....	..... ..... .....
Sorghum midge	..... .....	..... .....	..... ..... .....	..... ..... .....



Diagram of major pests of cereal crops:


Draw the life cycle of rice stem borer





Diagram of damaging symptoms of *Chiloptellus*



## EXERCISE NO. 3





### Objective: To identify insect pests attacking pulse crops

**Exercise:** Observe and enlist major pests attacking pulse crops. Draw the different life stages of *Helicoverpaarmigera*.

Common name	Scientific name	Family & Order	Pest identification/ Bionomics	Damaging symptoms
Gram pod borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Cut worm	..... .....	..... .....	..... ..... .....	..... ..... .....
Plume moth	..... .....	..... .....	..... ..... .....	..... ..... .....
Spotted pod borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Bean aphid	..... .....	..... .....	..... ..... .....	..... ..... .....

Leaf hopper	..... .....	..... .....	..... ..... .....	..... ..... .....
Thrips	..... .....	..... .....	..... ..... .....	..... ..... .....
Pod bug	..... .....	..... .....	..... ..... .....	..... ..... .....
Blister beetle	..... .....	..... .....	..... ..... .....	..... .....

**Life stages of *Helicoverpa armigera***



**EXERCISENO. 4**

**Objective: To identify insect pests attacking cash crops**

**Exercise:** Observe and enlist major pests attacking cash crops. Write the integrated pest management of sugarcane.

Common name	Scientific name	Family & Order	Pest identification/ Bionomics	Damaging symptoms
<b>Pests of Sugarcane</b>				
Early Shoot borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Internode borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Top borer	..... .....	..... .....	..... ..... .....	..... ..... .....
White grub	..... .....	..... .....	..... ..... .....	..... ..... .....
Termites	..... .....	..... .....	..... ..... .....	..... ..... .....

Pyrilla	..... .....	..... .....	..... ..... .....	..... ..... .....
Wooly aphid	..... .....	..... .....	..... ..... .....	..... ..... .....
<b>Pests of Cotton</b>				
Aphids	..... .....	..... .....	..... ..... .....	..... ..... .....
Whiteflies	..... .....	..... .....	..... ..... .....	..... ..... .....
Thrips	..... .....	..... .....	..... ..... .....	..... ..... .....
Red cotton bug	..... .....	..... .....	..... ..... .....	..... ..... .....

			.....	.....
American bollworm	..... .....	..... .....	..... ..... .....	..... ..... .....
Pink boll worm	..... .....	..... .....	..... ..... .....	..... ..... .....
Spiny bollworm	..... .....	..... .....	..... ..... .....	..... ..... .....
Tobacco caterpillar	..... .....	..... .....	..... ..... .....	..... ..... .....





**EXERCISE NO. 5**

**Objective: To identify insect pests attacking vegetable crops**

**Exercise:** Observe and enlist major pests attacking vegetable crops. Draw the diagram of the observed pests.

Common name	Scientific name	Family & Order	Pest identification/ Bionomics	Damaging symptoms
<b>Pests of Solanaceous crops</b>				
Fruit borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Tobacco caterpillar	..... .....	..... .....	..... ..... .....	..... ..... .....
White flies	..... .....	..... .....	..... ..... .....	..... ..... .....
Thrips	..... .....	..... .....	..... ..... .....	..... ..... .....
Pin worm	..... .....	..... .....	..... ..... .....	..... ..... .....

Serpentine leaf miner	..... .....	..... .....	..... ..... .....	..... ..... .....
Fruit sucking moth	..... .....	..... .....	..... ..... .....	..... ..... .....
Brinjal shoot and fruit borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Epilachna beetle	..... .....	..... .....	..... ..... .....	..... ..... .....
Chilli thrips	..... .....	..... .....	..... ..... .....	..... ..... .....
Aphids	..... .....	..... .....	..... ..... .....	..... ..... .....





Potato tuber moth	..... .....	..... .....	..... ..... .....	..... ..... .....
Cut worms	..... .....	..... .....	..... ..... .....	..... ..... .....
Potato GLH	..... .....	..... .....	..... ..... .....	..... ..... .....
White grubs	..... .....	..... .....	..... ..... .....	..... ..... .....
<b>Pests of Okra</b>				
Bhendi fruit and shoot borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Bhendi fruit borer	..... .....	..... .....	..... ..... .....	..... ..... .....

White fly	..... .....	..... .....	..... ..... .....	..... ..... .....
Aphid	..... .....	..... .....	..... ..... .....	..... ..... .....
Jassids	..... .....	..... .....	..... ..... .....	..... ..... .....
Red cotton bug	..... .....	..... .....	..... ..... .....	..... ..... .....
<b>Pest of Crucifers</b>				
Diamond back moth	..... .....	..... .....	..... ..... .....	..... ..... .....
Cabbage borer	..... .....	..... .....	..... ..... .....	..... ..... .....

Cabbage green semilooper	..... .....	..... .....	..... ..... .....	..... ..... .....
Cabbage butterfly	..... .....	..... .....	..... ..... .....	..... ..... .....
Tobacco caterpillar	..... .....	..... .....	..... ..... .....	..... ..... .....
Cabbage aphid	..... .....	..... .....	..... ..... .....	..... ..... .....
Mustard aphid	..... .....	..... .....	..... ..... .....	..... ..... .....
<b>Pests of Cucurbits</b>				
Fruit flies	..... .....	..... .....	..... ..... .....	..... ..... .....

Pumpkin beetles	..... .....	..... .....	..... ..... .....	..... ..... .....
Stem gall fly	..... .....	..... .....	..... ..... .....	..... ..... .....
Pumpkin caterpillar	..... .....	..... .....	..... ..... .....	..... ..... .....
Leaf miner	..... .....	..... .....	..... ..... .....	..... ..... .....

**Diagram of important pests of vegetable crops**





**EXERCISE NO. 6**


**Objective: To identify insect pests attacking oilseed crops**

**Exercise:** Observe and enlist major pests attacking oilseed crops.

Common name	Scientific name	Family & Order	Pest identification/ Bionomics	Damaging symptoms
Capitulum borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Bihar Hairy Caterpillar	..... .....	..... .....	..... ..... .....	..... ..... .....
Tobacco caterpillar	..... .....	..... .....	..... ..... .....	..... ..... .....
Red Hairy Caterpillar	..... .....	..... .....	..... ..... .....	..... ..... .....
Groundnut leaf miner	..... .....	..... .....	..... ..... .....	..... ..... .....

White grub	..... .....	..... .....	..... ..... .....	..... ..... .....
Mustard aphid	..... .....	..... .....	..... ..... .....	..... ..... .....
Mustard sawfly	..... .....	..... .....	..... ..... .....	..... ..... .....
Sesame leaf hopper	..... .....	..... .....	..... ..... .....	..... ..... .....
Sphingid	..... .....	..... .....	..... ..... .....	..... ..... .....


**Draw the neat diagram of the following pests**




**White grub**



**Aphid**



**Leaf miner symptoms**



**Springid**

**EXERCISE NO. 7**

**Objective: To identify insect pests attacking fruit crops**

**Exercise:** Observe and enlist major pests attacking fruit crops.

Common name	Scientific name	Family & Order	Pest identification/ Bionomics	Damaging symptoms
<b>Pests of Mango</b>				
Mango hopper	..... .....	..... .....	..... ..... .....	..... ..... .....
Stem borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Fruit fly	..... .....	..... .....	..... ..... .....	..... ..... .....
Mango nut weevil	..... .....	..... .....	..... ..... .....	..... ..... .....
Mango mealy bug	..... .....	..... .....	..... ..... .....	..... ..... .....

Bark eating caterpillar	..... .....	..... .....	..... ..... .....	..... ..... .....
		<b>Pest of Banana</b>		
Rhizome weevil	..... .....	..... .....	..... ..... .....	..... ..... .....
Pseudostem borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Banana aphid	..... .....	..... .....	..... ..... .....	..... ..... .....
		<b>Pest of apple, pear and plum</b>		
Apple woolly aphid	..... .....	..... .....	..... ..... .....	..... ..... .....
San Jose scale	..... .....	..... .....	..... ..... .....	..... ..... .....

Cotton cushiony scale	..... .....	..... .....	..... ..... .....	..... ..... .....
Codling moth	..... .....	..... .....	..... ..... .....	..... ..... .....
Green peach aphid	..... .....	..... .....	..... ..... .....	..... ..... .....
		<b>Pests of citrus</b>		
Shoot psyllid	..... .....	..... .....	..... ..... .....	..... ..... .....
Citrus leaf miner	..... .....	..... .....	..... ..... .....	..... ..... .....
Citrus whitefly	..... .....	..... .....	..... ..... .....	..... ..... .....
Fruit piercing moth	..... .....	..... .....	..... ..... .....	..... ..... .....

			.....	.....
Bark eating caterpillar	..... .....	..... .....	..... ..... .....	..... ..... .....
Citrus butterfly	..... .....	..... .....	..... ..... .....	..... ..... .....
		<b>Pests of guava</b>		
Fruit borer complex	..... .....	..... .....	..... ..... .....	..... ..... .....
Fruit flies	..... .....	..... .....	..... ..... .....	..... ..... .....
Bark borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Tailed mealy bug	..... .....	..... .....	..... ..... .....	..... ..... .....

Spiralling white fly	..... .....	..... .....	..... ..... .....	..... ..... ..... .....
		<b>Pests of pomegranate</b>		
Anar butterfly	..... .....	..... .....	..... ..... .....	..... ..... .....
Tailed mealy bug	..... .....	..... .....	..... ..... .....	..... ..... .....
Aphids	..... .....	..... .....	..... ..... .....	..... ..... .....
White fly	..... .....	..... .....	..... ..... .....	..... ..... .....
		<b>Pests of grapevine</b>		
Stem girdler	..... .....	..... .....	..... ..... .....	..... ..... .....



Flea beetle	..... .....	..... .....	..... ..... .....	..... ..... .....
Thrips	..... .....	..... .....	..... ..... .....	..... ..... .....
Mealy bugs	..... .....	..... .....	..... ..... .....	..... ..... .....

**EXERCISE NO. 8**

**Objective: To identify insect pests attacking plantation crops**

**Exercise:** Observe and enlist major pests attacking plantation crops. Write the management practices of rhinoceros beetle attacking coconut.

Common name	Scientific name	Family & Order	Pest identification/ Bionomics	Damaging symptoms
Rhinoceros beetle	..... .....	..... .....	..... ..... .....	..... ..... .....
Red palm weevil	..... .....	..... .....	..... ..... .....	..... ..... .....
Spindle bug (oilpalm/ arecanut)	..... .....	..... .....	..... ..... .....	..... ..... .....
Termites	..... .....	..... .....	..... ..... .....	..... ..... .....
Root grubs	..... .....	..... .....	..... ..... .....	..... ..... .....

Arecanut Inflorescence caterpillar	..... .....	..... .....	..... ..... .....	..... ..... .....
Mites	..... .....	..... .....	..... ..... .....	..... ..... .....
Coffee berry borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Mealy bugs	..... .....	..... .....	..... ..... .....	..... ..... .....
Tea mosquito bug	..... .....	..... .....	..... ..... .....	..... ..... .....
Tea mite complex	..... .....	..... .....	..... ..... .....	..... ..... .....



**EXERCISE NO. 9**

**Objective: To identify insect pests attacking spices, condiments, and narcotics**

**Exercise:** Observe and enlist major insect pests attacking spices, condiments, and narcotics.

Common name	Scientific name	Family & Order	Pest identification/ Bionomics	Damaging symptoms
<b>Pests of spices and condiments</b>				
Cardamom thrips	..... .....	..... .....	..... ..... .....	..... ..... .....
Cardamom Aphid	..... .....	..... .....	..... ..... .....	..... ..... .....
Cardamom capsule borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Cardamom hairy caterpillar	..... .....	..... .....	..... ..... .....	..... ..... .....
Pepper pollu beetle	..... .....	..... .....	..... ..... .....	..... ..... .....
Tailed mealy bug	..... .....	..... .....	..... ..... .....	..... ..... .....

Ginger shoot borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Ginger Rhizome scale	..... .....	..... .....	..... ..... .....	..... ..... .....
Thrips	..... .....	..... .....	..... ..... .....	..... ..... .....
		<b>Pest of tobacco</b>		
Tobacco caterpillar	..... .....	..... .....	..... ..... .....	..... ..... .....
Tobacco stem borer	..... .....	..... .....	..... ..... .....	..... ..... .....
Whitefly	..... .....	..... .....	..... ..... .....	..... ..... .....
Aphid	..... .....	..... .....	..... ..... .....	..... ..... .....

**Objective: To identify insect pests attacking ornamental plants**


**Exercise:** Observe and enlist major insect pests attacking ornamental plants. Draw the different life stages of AK butterfly.

Common name	Scientific name	Family & Order	Pest identification/ Bionomics	Damaging symptoms
Rose thrips	..... .....	..... .....	..... ..... .....	..... ..... .....
Rose aphids	..... .....	..... .....	..... ..... .....	..... ..... .....
Leaf cutter bee	..... .....	..... .....	..... ..... .....	..... ..... .....
Dusky cotton bug	..... .....	..... .....	..... ..... .....	..... ..... .....
Banded blister beetle	..... .....	..... .....	..... ..... .....	..... ..... .....
Ak butterfly	..... .....	..... .....	..... ..... .....	..... ..... .....


Lily moth	..... .....	..... .....	..... ..... .....	..... ..... .....
Gerbera leaf miner	..... .....	..... .....	..... ..... .....	..... ..... .....




**Life stages of AK butterfly**




**Egg**



**Larva**



**Pupa**



**Adult**

**EXERCISE NO. 11**

**Objective: To identify insect pests and Mites associated with stored grain**

**Exercise:** Observe and enlist the insect pests and mites attacking stored grain

Common name	Scientific name	Family & Order	Pest identification/ Bionomics	Damaging symptoms
Rice weevil	..... .....	..... .....	..... ..... .....	..... ..... .....
Angoumois grain moth	..... .....	..... .....	..... ..... .....	..... ..... .....
Cigarette beetle	..... .....	..... .....	..... ..... .....	..... ..... .....
Red flour beetle	..... .....	..... .....	..... ..... .....	..... ..... .....
Rice moth	..... .....	..... .....	..... ..... .....	..... ..... .....

Khapra beetle	..... .....	..... .....	..... ..... .....	..... ..... .....
Saw toothed grain beetle	..... .....	..... .....	..... ..... .....	..... ..... .....
Long headed flour beetle	..... .....	..... .....	..... ..... .....	..... ..... .....
Grain mite	..... .....	..... .....	..... ..... .....	..... ..... .....

**Objective: To determine insect infestation**

**Exercise: Explain various methods for determination of insect**

**Grain Probe Traps:** .....

.....

.....

.....

.....

.....

**Sticky Traps:**.....

.....

.....

.....

.....

.....

**Refuse Trap Method:**.....

.....

.....

.....

.....

.....

.....

.....

**Light Trap Method**.....

.....

.....

.....

.....

.....

.....

.....

**Use of Pheromone**.....

.....

.....

.....

.....  
.....

**Visual Lures:**.....

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

**Acoustical Methods:**.....

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

**Electrical Conductance:**.....

.....  
.....  
.....  
.....  
.....

**Berlese Funnel Method:**.....

.....  
.....  
.....

**X-Ray Imaging:** .....

.....  
.....  
.....  
.....

**Objective: To assess losses due to insects**

**Exercise:** Visit to field and asses the losses caused by the insect

**Basic needs:**

.....

.....

.....

.....

.....

.....

**Assessment of Losses due to Insect Pests**













**Objective: To study non-insect pests in field/godown**

**Exercise:** Identify the non- insect pests attacking in field/ godown and write their management

**Observation:**

1. Giant African snail:.....  
.....  
.....

2. Millipedes:.....  
.....  
.....

3. Mites:.....  
.....  
.....

4. Snails and slugs:.....  
.....  
.....

5. Nematodes:.....  
.....  
.....

6. Rodents: .....  
.....  
.....

7. Aves.....  
.....  
.....

Others: .....  
.....  
.....

.....  
.....





.....  
.....

**Observations**

<b>S.No.</b>	<b>Item</b>	<b>Sample no.</b>	<b>Weight</b>
1	Initial weight of sample	1	
		2	
		3	
2	Fresh weight of sample	1	
		2	
		3	
3	Container weight without sample	1	
		2	
		3	

**Results:** Tabulate the results as follows:

<b>S. No.</b>	<b>Moisture measurement method</b>	<b>Sample No. (wet basis)</b>	<b>Moisture content (dry basis)</b>	<b>Moisture content</b>
1	Air oven method	1		
		2		
		3		
2	Universal moisture meter method	1		
		2		
		3		
3	Brown dual fractional distillation method	1		
		2		
		3		







## APPENDICES

### GENERAL SYMPTOMS OF DAMAGE CAUSED BY DIFFERENT PESTS ON CROP PLANTS

Entomologist classified the plant feeding insect into two categories such as generalist and specialist according to the mode of host plant use by them. Generalist insect can be defined those insect which use wide range of plant species as their host, whereas the specialist insect using a specified range of host plants in their life stages. Again phytophagous insects are differentiated into three categories such as *monophagous*, *oligophagous* and *polyphagous*. The insect species which feed on plants under single genus termed as monophagous. The oligophagous type consumed wide range of plants of different genera but in a single plant family. Whereas a polyphagous insect refers that they are feeding wide range of plant under different plant families. Most of the phytophagous insects are specialized for choosing their host plant.

**Root feeding insects:** Insect larvae feed on roots, root nodules; nymphs and adults suck sap from roots, resulting in stunted growth, poor tillering, drying of plants in isolated patches e.g: White grubs, grubs of rhinoceros beetles, termites, rice root weevil and ragi root aphid

**Stem borers:** Larvae enter into the shoot of main stem, tillers and feed on the central growing point. As a result, nutrient supply from the main plant beyond the infested part is affected leading to withering, wilting and drying up exhibiting symptoms such as dead heart / white ear / over growths of bunchy tops. Stem borers of paddy, millets, sugarcane, brinjal, bhendi, cotton etc.

**Shoot borers:** Larvae attack tender shoots and bore inside during the vegetative stage of crop growth and cause wilting, dropping of terminal plant part which later dries up e.g., Shoot fly of sorghum, early stem borer in sugarcane, stem fly in black gram/French bean, soybean, shoot borers of brinjal, bhendi, cotton, castor etc.

**Tree borers:** Larvae bore deep into the tree trunk, tunnels in zigzag manner and feed on inner tissues, affecting nutrient and translocation of sap to upper portions of branches / tree exhibiting symptoms such as withering of leaves, drying of twigs or complete dying of tree. Presence of fresh powdered material, ooze of gummy exudations etc. from the affected portion on the tree trunk is also seen in some cases e.g., Tree borers of mango, cashew; coconut red palm weevil etc.

**Bark borers:** Larvae enter into the bark and tunnel into the branches. The larvae remain hiding in the galleries formed from floss / fecal matter and silken saliva on the stem and continue to scrape the bark. Larval feeding results in drying of branches and breaking of affected portion with wind or gale e.g., Bark eating caterpillars of citrus, mango, guava, casuarinas, jack etc.

**Gall formers:** Larvae/nymphs feeding inside the stem/ tiller /leaf/ flower bud affect the tissue by nibbling the meristematic tissues and secretion of auxins that results in excessive growth of cells at the affected portion leading to distorted growth and malformation of plant parts known as 'Gall' e.g., Paddy gall midge, chilly midge, gingelly midge, cucurbit stem borer, mango malformations, tobacco stem borer, cotton stem weevil, mango inflorescence midge, chilli midge etc.

**Leaf folders:** Larvae fasten the margins of individual leaves from margins / fold longitudinally or roll leaves into bell shape and feeds within by scrapping the chlorophyll e.g., Rice/ maize/leaf folder, cotton leaf roller, red gram/ black gram leaf folder.

**Leaf miners:** Larvae fasten the leaves /leaflets by means of silken threads (derived from saliva) and scrape the chlorophyll content by remaining within the web. Fecal pellets / frass remains present in the web e.g., Leaf webbers on groundnut / gingelly, Webbers of mango / sapota /Cashew.

**Leaf webbers:** Larvae fasten the leaves /leaflets by means of silken threads (derived from saliva) and scrape the chlorophyll content by remaining within the web. Fecal pellets / frass remains present in the web e.g., Leaf webbers on groundnut / gingelly, Webbers of mango / sapota /Cashew.

**Defoliators / Skeletonizers:** Larvae feed on the leaves completely leaving only midrib / veins or scrape the chlorophyll content of leaves or cause numerous holes e.g., Castor semilooper, red hairy caterpillar, Bihar hairy caterpillar, snakegourd semilooper, ash weevils, tobacco caterpillar, brinjal epilachna beetle.

**Pod / Capsule borers:** During the reproductive stage of crop, larvae bore into the flowers, pods, capsules and fasten the adolescent plant parts with silken threads, frass and excreta and feed on the internal contents within the web e.g., Spotted pod borer in legumes, capsule borers of castor / gingelly ; pod borer complex in pulses, gram caterpillar, pink bollworm, tobacco caterpillar , chilly pod borer etc.

**Fruit borers / Bollworms:** Larvae enter into the tender fruit's bolls and feed on internal content /pulp and plug the larval burrow with excreta e.g., Fruit borer of brinjal /bhendi /tomato, mango fruit borer, fruit fly, mango stone weevil, cashew apple and nut borer, anar/ guava fruit borer, Cotton bollworm etc.

**Seed feeding insects / stored grain pests:** Larvae feed on stored seeds either as internal/ external feeders / by webbing the food particles e.g., Rice weevil, lesser grain borer, red rust flour beetle, rice moth, cigarette beetle, saw toothed beetle.

**Sap sucking insects / feeders:**

**From tender plant parts:** Nymphs and adults suck sap from the base of the plant/leaves / tender terminal plant parts and affect the vigor and growth of the plants. Different insects exhibit different symptoms. Most of the sap suckers suck sap in excess of their requirement and excrete honey dew, which is rich in sugars, a source for sooty mold development e.g., Aphids, leafhoppers (jassids), plant hoppers, white flies etc. on important crops.


**From grains:** Nymphs and adults suck juice from developing ovaries/milky grains resulting in the formation of shriveled /chaffy grains e.g., Rice gundhy bug, sorghum ear-head bug, sorghum midge.

**Symptoms of sucking pest damage:**

- Hopper burn (drying of margins), complete drying of plants in patches - paddy brown plant hopper, cotton leafhoppers
- Scorch appearance of leaves - paddy leafhopper
- Upward curling of leaves - Chilli thrips
- Downward curling of leaves and elongation of petioles of older leaves-Chilli white mites
- Leaf drying from tip down wards - Onion thrips
- Yellowing /crinkling of leaves - Thrips on groundnut, pulses
- White / yellow blotches on upper surface of leaves - Thrips in castor, Cotton mites, Brinjal mites
- Red streaks on leaves - Sorghum/Maize mites
- Reduced vigour /sooty mold /yellowing / - Aphids / whiteflies on cotton/brinjal
- Corky out growths on fruits - tea mosquito bug in guava
- Rotting of fruits - fruit sucking moth in citrus fruit flies in mango
- Die back symptoms - tea mosquito bug in cashew

## MAJOR INSECT PESTS ATTACKING CEREAL CROPS

Common name	Scientific name	Family	Order
<b>Paddy</b>			
Rice yellow stem borer	<i>Scirpophagaincertulas</i>	Pyraustidae	Lepidoptera
Gall midge	<i>Orseoliaoryzae</i>	Cecidomyiidae	Diptera
Green leaf hopper	<i>Nephotettixvirescens</i> , <i>N. nigropictus</i> ; <i>N. cincticeps</i>	Cicadellidae	Hemiptera
Brown plant hopper	<i>Nilaparvatalugens</i>	Delphacidae	Hemiptera
White back plant hopper	<i>Sogatellafurcifera</i>	Delphacidae	Hemiptera
Rice earhead bug	<i>Leptocorisa acuta</i>	Alydidae	Hemiptera
Rice leaf folder	<i>Cnaphalocrocismedinalis</i>	Pyralidae	Lepidoptera
Rice caseworm	<i>Nymphuladepunctalis</i>	Pyraustidae	Lepidoptera
Rice grasshopper	<i>Hieroglyphus banian</i>	Acrididae	Orthoptera
<b>Wheat</b>			
Wheat aphid	<i>Macrosiphum miscanthi</i>	Aphididae	Hemiptera
Climbing cutworm/armyworm	<i>Mythimna separata</i>	Noctuidae	Lepidoptera
Ghujhia Weevil	<i>Tanymecus indicus</i>	Curculionidae	Coleoptera
Termites	<i>Odontotermesobesus/ Microtermesobesi</i>	Termitidae	Isoptera
<b>Maize and Sorghum</b>			
Stem borer	<i>Chilopartellus</i>	Crambidae	Lepidoptera
Fall armyworm	<i>Spodoptera frugiperda</i>	Noctuidae	Lepidoptera
Shoot fly	<i>Atherigonasoccata; Atherigonaorientalis</i>	Muscidae	Diptera
Shoot bug	<i>Peregrinus maidis</i>	Delphacidae	Hemiptera
Sorghum midge	<i>Contariniatorghicola</i>	Cecidomyiidae	Diptera



**Rice stem borer**

**Green leaf hopper**

**Rice leaf folder**



**Termites**



© RAYT BERTONE 2014


**Fall armyworm larva**



**Shootfly**

## INSECT PESTS ATTACKING PULSE CROPS

Common name	Scientific name	Family	Order
Gram pod borer	<i>Helicoverpaarmigera</i>	Noctuidae	Lepidoptera
Plume moth	<i>Exelastisatomosa</i>	Pterophoridae	Lepidoptera
Spotted pod borer	<i>Marucatestulalis</i>	Pyraustidae	Lepidoptera
Bean aphid	<i>Aphis craccivora</i>	Aphididae	Hemiptera
Leaf hopper	<i>Empoascaferri, E. binotata, E. flavescens</i>	Cicadellidae	Hemiptera
Thrips	<i>Ayyariachaetophora, Caliothrips indicus, Megalurothrips distalis</i>	Thripidae	Thysanoptera
Pod bug	<i>Riptortuspedestris</i>	Coreidae	Hemiptera
Blister beetle	<i>Mylabrispustulata</i>	Meloidae	Coleoptera




Life stages of *Helicoverpaarmigera*

Damaging stage of blister beetle

## INSECT PESTS ATTACKING CASH CROPS

Common name	Scientific name	Family	Order
<b>Sugarcane</b>			
Early Shoot borer	<i>Chiloinfuscatellus</i>	Crambidae	Lepidoptera
Internode borer	<i>Chilosacchariphagusindicus</i>	Crambidae	Lepidoptera
Top borer	<i>Scirpophagaexcerptalis</i>	Pyralidae	Lepidoptera
White grub	<i>Holotrichiaconsanguinea</i>	Melolonthidae	Coleoptera
Termites	<i>Odontotermesobesus</i>	Termitidae	Isoptera
Pyrilla	<i>Pyrillaperpusilla</i>	Lophopidae	Hemiptera
Woolly aphid	<i>Ceratovacunalanigera</i>	Aphididae	Hemiptera
<b>Cotton</b>			
Aphids	<i>Aphis gossypii</i>	Aphididae	Hemiptera
Whiteflies	<i>Bemisiatabaci</i>	Aleyrodidae	Hemiptera
Thrips	<i>Thrips tabaci</i>	Thripidae	Thysanoptera
Red cotton bug	<i>Dysdercuscingulatus</i>	Pyrrhocoridae	Hemiptera
American bollworm	<i>Helicoverpaarmigera</i>	Noctuidae	Lepidoptera
Pink boll worm	<i>Pectinophoragossypiella</i>	Gelechiidae	Lepidoptera
Spotted boll worm	<i>Eariasvittella &amp; E. insulana</i>	Noctuidae	Lepidoptera





### INSECT PESTS ATTACKING VEGETABLE CROPS

Common name	Scientific name	Family	Order
<b>Solanaceous crops</b>			
Fruit borer	<i>Helicoverpaarmigera</i>	Noctuidae	Lepidoptera
Tobacco caterpillar	<i>Spodoptera litura</i>	Noctuidae	Lepidoptera
White flies	<i>Bemisiatabaci</i>	Aleyrodidae	Hemiptera
Thrips	<i>T. tabaci, F. schultzi</i>	Thripidae	Thysanoptera
Pin worm	<i>Tuta absoluta</i>	Gelechiidae	Lepidoptera
Serpentine leaf miner	<i>Liriomyzatrifolii</i>	Agromyzidae	Diptera
Fruit sucking moth	<i>Othreisfullonica, O. materna, O. ancilla</i>	Noctuidae	Lepidoptera
Brinjal shoot and fruit borer	<i>Leucinodes orbonalis</i>	Pyraustidae	Lepidoptera
Epilachna beetle	<i>Henosepilachnadodecastigma, H. vigintioctopunctata, H. demurille, H. implicata</i>	Coccinellidae	Coleoptera
Chilli thrips	<i>Scirtothrips dorsalis</i>	Thripidae	Thysanoptera
Aphids	<i>Myzuspersicae</i>	Aphididae	Hemiptera
Potato tuber moth	<i>Phthorimaeaoperculella</i>	Gelechiidae	Lepidoptera
Cut worms	<i>Agrotisipsilon, A. segetum, Xestia C. nigrum and Peridromasauca</i>	Noctuidae	Lepidoptera
Potato GLH	<i>Empoascakerri</i>	Cicadellidae	Hemiptera
White grubs	<i>Holotrichiaexcisa, H. repetita, H. notaticollis, Anomala communis, A. nathani</i>	Melolonthidae	Coleoptera
<b>Okra</b>			
Bhendi shoot and fruit borer	<i>Eariasvitella, E. insulana</i>	Noctuidae	Lepidoptera
Bhendi fruit borer	<i>Helicoverpaarmigera</i>	Noctuidae	Lepidoptera
White fly	<i>Bemisiatabaci</i>	Aleyrodidae	Hemiptera
Aphid	<i>Aphis gossypii</i>	Aphididae	Hemiptera




Jassids	<i>Amrascabigutulabigutula</i>	Cicadellidae	Hemiptera
Red cotton bug	<i>Dysdercuskoenigii</i>	Pyrrhocoridae	Hemiptera
<b>Crucifers</b>			
Diamond back moth	<i>Plutellaxylostella</i>	Plutellidae	Lepidoptera
Cabbage borer	<i>Hellulaundalis</i>	Pyraustidae	Lepidoptera
Cabbage green semilooper	<i>Tircihoplusiani</i>	Noctuidae	Lepidoptera
Cabbage butterfly	<i>Pieris brassicae</i>	Pieridae	Lepidoptera
Tobacco caterpillar	<i>Spodoptera litura</i>	Noctuidae	Lepidoptera
Cabbage aphid	<i>Brevicorynebrassicae</i>	Aphididae	Hemiptera
Mustard aphid	<i>Lipaphiserysimi</i>	Aphdidae	Hemiptera
<b>Cucurbits</b>			
Fruit flies	<i>Bactroceracucurbitae</i>	Tephritidae	Diptera
Pumpkin beetles	<i>Aulacophorafoveicollis, A. cincta, A. intermedia</i>	Galerucidae	Coleoptera
Stem gall fly	<i>Neolasoipterafalcate</i>	Cecidomyiidae	Diptera
Pumpkin caterpillar	<i>Plusiapeponis, P. signata, P. orichalcea</i>	Noctuidae	Lepidoptera
Leaf miner	<i>Liriomyzatrifolii</i>	Agromyzidae	Diptera



**INSECT PESTS ATTACKING OILSEED CROPS**










Common name	Scientific name	Family	Order
Capitulum borer	<i>Helicoverpaarmigera</i>	Noctuidae	Lepidoptera
Bihar Hairy Caterpillar	<i>Spilosomaobliqua</i>	Arctiidae	Lepidoptera
Tobacco caterpillar	<i>Spodoptera litura</i>	Noctuidae	Lepidoptera
Red Hairy Caterpillar	<i>Amsactaalbistriga</i>	Arctiidae	Lepidoptera
Groundnut leaf miner	<i>Aproaeremamodicella</i>	Gelechiidae	Lepidoptera
White grub	<i>Holotrichiaconsanguinea</i>	Melolonthidae	Coleoptera
Mustard aphid	<i>Lipaphiserysimi</i>	Aphididae	Hemiptera
Mustard sawfly	<i>Athalialugens</i>	Tenthredinidae	Hymenoptera
Sesame leaf hopper	<i>Orosiusalbicinctus</i>	Cicadallidae	Hemiptera
Sphingid	<i>Acherontiastryx</i>	Sphingidae	Lepidoptera



### PESTS ATTACKING FRUIT CROPS

Common name	Scientific name	Family	Order
<b>Mango</b>			
Mango hopper	<i>Idioscopusniveosparsus</i> , <i>l. clypealis</i> , <i>Amritodusatkinsoni</i>	Cicadellidae	Hemiptera
Stem borer	<i>Batocerarufomaculata</i>	Cerambycidae	Coleoptera
Fruit fly	<i>Bactrocera dorsalis</i>	Tephritidae	Diptera
Mango nut weevil	<i>Sternochaetusmangiferae</i>	Curculionidae	Coleoptera
Mango mealy bug	<i>Drosichamangiferae</i>	Margarodidae	Hemiptera
Bark eating caterpillar	<i>Indarbelatetraonis</i> , <i>l. quadrinotata</i>	Metarbelidae	Lepidoptera
<b>Banana</b>			
Rhizome weevil	<i>Cosmopolites sordidus</i>	Curculionidae	Coleoptera
Pseudostem borer	<i>Odoiporuslongicollis</i>	Curculionidae	Coleoptera
Banana aphid	<i>Pentalonianigronevosa</i>	Aphididae	Hemiptera
<b>Apple, pear and plum</b>			
Apple woolly aphid	<i>Eriosomalanigerum</i>	Pemphigidae	Hemiptera
San Jose scale	<i>Quadraspidiotusperniciosus</i>	Diaspididae	Hemiptera
Cotton cushiony scale	<i>Iceryapurchase</i>	Margarodidae	Hemiptera
Codling moth	<i>Cydiaopomonella</i>	Tortricidae	Lepidoptera
Green peach aphid	<i>Myzuspersicae</i>	Aphididae	Hemiptera
<b>Citrus</b>			
Shoot psyllid	<i>Diaphorinacitri</i>	Psyllidae	Hemiptera
Citrus leaf miner	<i>Phyllocnistiscitrella</i>	Gracillariidae	Lepidoptera
Citrus whitefly	<i>Dialeurodescitri</i>	Aleyrodidae	Hemiptera


Fruit piercing moth	<i>Othreis materna, O. fullonica, O. ancilla</i>	Noctuidae	Lepidoptera
Bark eating caterpillar	<i>Indarbelatetraonis</i>	Metarbelidae	Lepidoptera
Citrus butterfly	<i>Papiliodemoleus, P. polytes</i>	Papilionidae	Lepidoptera
<b>Guava</b>			
Fruit borer	<i>Virachola (Duodorix) isocrates</i>	Lycaenidae	Lepidoptera
Fruit flies	<i>Bactrocera (Dacus) diversus</i>	Tephritidae	Diptera
Bark borer	<i>Indarbelatetraonis</i>	Metarbelidae	Lepidoptera
Tailed mealy bug	<i>Ferrisia virgata, Maconellicoccus hirsutus</i>	Pseudococcidae	Hemiptera
Spiralling white fly	<i>Aleurodicus disperses</i>	Aleyrodidae	Hemiptera
<b>Pomegranate</b>			
Anar butterfly	<i>Duodorix Isocrates</i>	Lycaenidae	Lepidoptera
Tailed mealy bug	<i>Ferrisia virgata, Maconellicoccus hirsutus</i>	Pseudococcidae	Hemiptera
Aphids	<i>Aphis punicae</i>	Aphididae	Hemiptera
White fly	<i>Aleurodicus disperses</i>	Aleyrodidae	Hemiptera
<b>Grapevine</b>			
Stem girdler	<i>Sthenias grisator</i>	Cerambycidae	Coleoptera
Flea beetle	<i>Scelodonta strigicollis</i>	Eumolpidae	Coleoptera
Thrips	<i>Rhipiphoro thrips cruentatus</i>	Thripidae	Thysanoptera
Mealy bugs	<i>Maconellicoccus hirsutus</i>	Pseudococcidae	Hemiptera

			
<b>Mango hopper</b>	<b>Mango stem borer</b>	<b>Mango nut weevil</b>	
			
<b>Leaf mining symptoms</b>	<b>Citrus butterfly</b>	<b>Stem girdler</b>	<b>Banana Pseudostem weevil symptoms</b>

### INSECT PESTS ATTACKING PLANTATION CROPS

Common name	Scientific name	Family	Order
Rhinoceros beetle	<i>Oryctes rhinoceros</i>	Scarabaeidae	Coleoptera
Red palm weevil	<i>Rhynchophorus ferrugineus</i>	Curculionidae	Coleoptera
Spindle bug	<i>Carvalhoia arecae</i>	Miridae	Hemiptera
Termites	<i>Odontotermus obesus</i>	Termitidae	Isoptera
Root grubs	<i>Leucopholis burmeisteri</i>	Melolonthidae	Coleoptera
Inflorescence caterpillar	<i>Tirathabamundella</i>	Pyralidae	Lepidoptera
Mites	<i>Oligonychus indicus</i>	Tetranychidae	Acari

Coffee berry borer	<i>Hypothenemus hampei</i>	Scolytidae	Coleoptera
Mealy bugs	<i>Ferrisia virgata</i> , <i>Planococcus lilacinus</i> , <i>P. citri</i>	Pseudococcidae	Hemiptera
Tea mosquito bug	<i>Helopeltis theivora</i>	Miridae	Hemiptera
Red spider mite	<i>Oligonychus coffeae</i>	Tetranychidae	Acari
Pink mite	<i>Acaphylla theae</i>	Eriophyidae	Acari
Yellow mite	<i>Polyphagotarsonemus latus</i>	Tarsonemidae	Acari



### INSECT PESTS ATTACKING SPICES, CONDIMENTS AND NARCOTICS

Common name	Scientific name	Family	Order
<b>Spices and condiments</b>			
Cardamom thrips	<i>Sciothrips cardamom</i>	Thripidae	Thysanoptera
Cardamom Aphid	<i>Pentalonia nigronervosa</i>	Aphididae	Hemiptera
Cardamom capsule borer	<i>Dichocrocis punctiferalis</i>	Pyraustidae	Lepidoptera
Cardamom hairycaterpillar	<i>Eupterotecardamomi</i>	Bombycidae	Lepidoptera
Pepper pollen beetle	<i>Longitarsus nigripennis</i>	Alticidae	Coleoptera
Ginger shoot borer	<i>Conogethes punctiferalis</i>	Pyraustidae	Lepidoptera
Rhizome scale	<i>Aspidiotus hartii</i>	Diaspididae	Hemiptera
Thrips	<i>Panchaetothrips indicus</i>	Thripidae	Thysanoptera
<b>Tobacco</b>			
Tobacco caterpillar	<i>Spodoptera litura</i>	Noctuidae	Lepidoptera
Stemborer	<i>Scrobipalpa heliopa</i>	Gelechiidae	Lepidoptera
Whitefly	<i>Bemisia tabaci</i>	Aleyrodidae	Hemiptera
Aphid	<i>Myzus nicotianae</i> , <i>Myzus persicae</i>	Aphididae	Hemiptera









### INSECT PESTS ATTACKING ORNAMENTAL PLANTS

Common name	Scientific name	Family	Order
Rose thrips	<i>Rhipiphorothripscruentatus</i>	Thripidae	Thysanoptera
Rose aphids	<i>Macrosiphumrosaeformis, M. rosae</i>	Aphididae	Hemiptera
Leaf cutter bee	<i>Megachileanthracina</i>	Megachilidae	Hymenoptera
Dusky cotton bug	<i>Oxycarenuslaetus</i>	Lygaeidae	Hemiptera
Banded blister beetle	<i>Mylabrisphalerata</i>	Meloidae	Coleoptera
Ak butterfly	<i>Danaischrysippus</i>	Nymphalidae	Lepidoptera
Lily moth	<i>Polytelagloriosae</i>	Noctuidae	Lepidoptera
Gerbera leaf miner	<i>Liriomyzatrifolii</i>	Agromyzidae	Diptera

		
<b><i>Danaischrysippus</i>(Larva)</b>	<b><i>Danaischrysippus</i>(Adult)</b>	<b>Leaf cutter bee</b>

### INSECT PESTS AND MITES ASSOCIATED WITH STORED GRAIN

Common name	Scientific name	Family	Order
Rice weevil	<i>Sitophilus oryzae,</i>	Curculionidae	Coleoptera
Angoumois grain moth	<i>Sitotrogacerealella</i>	Gelechiidae	Lepidoptera
Cigarette beetle	<i>Lasiodermasericorne</i>	Anobiidae	Coleoptera
Red flour beetle	<i>Triboliumcastaneum, T. confusum</i>	Tenebrionidae	Coleoptera
Rice moth	<i>Corcyra cephalonica</i>	Galleriidae	Lepidoptera
Khapra beetle	<i>Trogoderma granarium</i>	Dermestidae	Coleoptera
Saw toothed grain beetle	<i>Oryzaephillissurinamensis</i>	Silvanidae	Coleoptera
Long headed flour beetle	<i>Latheticusoryzae</i>	Tenebrionidae	Coleoptera
Grain mite	<i>Acarus siro</i>	Acaridae	Sarcoptiformes

		
<b>Rice weevil</b>	<b>Angoumois grain moth</b>	<b>Red flour beetle</b>
		
<b>Khapra beetle</b>	<b>Saw toothed grain beetle</b>	<b>Rice moth</b>

### DETERMINATION OF INSECT INFESTATION BY DIFFERENT METHODS

**Grain Probe Traps:** Grain probe traps are cylindrical tubes with perforations in the upper section through which insects drop into the trap and are unable to escape because of the shape of the receptacle. They are labour intensive, limits temporal availability of data, restricts placement of probe traps in easily accessible locations and difficult to interpret.

**Sticky Traps:** Any surface coated with a sticky substance (such as petroleum jelly or polybutene gel usually sold as bird repellent) that prevents insects from leaving after landing on it. They should be suspended from the store roof, to hang above or between stacks or heaps of stored grains. They have a short effective life since their surfaces are easily covered with dust.

**Refuse Trap Method:** These are made from waste material such as cardboard packaging. They provide a refuge for insects such as moth larvae which habitually leave the food source to pupate in sacking or crevices in the storage structure.

**Light Trap Method:** Most efficient at detecting moth infestations since the adults are attracted to light when they leave the produce in order to fly and mate. Ultraviolet (300-400nm) and green light (500-550nm) are the most attractive wavelengths to storage pests.

**Use of Pheromone:** For use in monitoring, chemical attractants are impregnated or encased in a rubber, paper or plastic lure that slowly releases the active components over a period of several days/weeks. Environmental factors affect catches: temperature, rainfall, wind speed and direction influence attractant release from lures and insect flight.

**Visual Lures:** Similar to light traps, visual lures are either lights that attract insects from the dark/dimly lit surroundings (usually fluorescent, incandescent and ultraviolet lights) or they are colored objects that are attractive because of their specific reflectance and shapes that stand out against a contrasting background. Electrical cutters are placed in dimly lit areas where their light is not visible outdoors such that it does not lure insects into the building.

**Acoustical Methods:** Use sound (insect feeding sounds) to automatically monitor both internal/external grain feeding insects. Affected by background noise, insect behaviour, insect inactivity, unfavorable environment, intensity/duration/spectral characteristics of the sound at source, distance of the receiver and receiver's spectral sensitivity.

**Electrical Conductance:** Conductance is monitored by measuring the voltage across the kernel (the kernel acts as one resistor in a two-resistor and voltage-divider circuit of the single kernel characterization system. This method requires skilled person.

**Berlese Funnel Method:** Works on the principle that insects move away from heat. It takes 5-6hrs to determine the presence of insects in 1Kg grain samples though proved 59% efficient in recovering Stephens adults in wheat. It is slow and inaccurate in detecting infestations; by this time, the grain would have been loaded into ships/bins.

**Near-Infrared Reflectance Spectroscopy:** Based on the absorption of electromagnetic wavelengths in the range of 780-2500nm to determine the concentrations of constituents like water, proteins, fats and carbohydrates using classical absorption spectroscopy. It has proved in several Coleopteran species and external and internal infestations in wheat, with up to 1000 kernels scan per second.

**Machine Vision:** Individual grain kernels are compared with the photographic print of the representative sample. It consists of high speed integrated machine vision software used with a monochrome CCD camera and a personal computer. It has a limited rate of sample throughout put

**X-Ray Imaging:** A soft X-ray system consisting of a fluoroscope operated at 15KV potential and 65 $\mu$  A, produces the real time non-destructive, highly accurate images. It can detect both internal and external insects, and able to detect both live and dead insects inside the grain kernels, except it can't detect insect eggs.

## ASSESSMENT OF LOSSES DUE TO INSECTS

### Basic Needs

- To determine the economic status of a given pest species.
- For establishing the economic threshold levels and economic injury levels of the pest.

- To estimate the effectiveness of control measures.
- For evaluating the crop or a variety for its reaction to the pests.
- Helping in deciding the allocations for research and extension in plant protection.
- Helping in assigning the priorities on the bases of relative importance of different pests.

### Assessment of Losses due to Insect Pests

**Stem borer:** Based on eggs and larval damage: Presence of yellowish-brown egg mass near the leaf tip/presence of dead heart (vegetative stage) or white ear (reproductive stage).

a. Eggs in the nursery: Number of egg masses/m<sup>2</sup> (ETL: 2)

b. Larval damage: Count the total tillers and affected tillers in a unit area and arrive at a percentage

$$\% \text{ drying branches} = \frac{\text{Number of drying branches}}{\text{Total number of branches}} \times 100 \text{ (ETL: 10\%)}$$

### Inflorescence midge:

$$\% \text{ Infected shoot} = \frac{\text{Number of infected bud}}{\text{Total number of bud}} \times 100 \text{ (ETL 10\%)}$$

**Leaf Webber:** Based on damage - folded and scrapped leaves

$$\% \text{ leaf damage} = \frac{\text{Number of damaged leaves}}{\text{Total number of leaves}} \times 100$$

(ETL: 10% at vegetative stage or 5% at flowering stage) (in 10 randomly selected plants)

**Whorl maggot:** Based on damage - marginal blotching and yellow patches on the leaves

$$\% \text{ leaf damage} = \frac{\text{Number of Damage Leaf}}{\text{Total number of leaves}} \times 100 \quad \text{(in 10 randomly selected plants)}$$

$$\% \text{ Avoidable yield losses} = \frac{\text{Yield in protected crop} - \text{Yield in unprotected crop}}{\text{Yield in protected crop}} \times 100$$

### PESTICIDE DOSAGE CALCULATION

The following formulas are useful in quantifying insecticides for field application.

1. For spraying

Preparation of spray solution is  $V_1 S_1 = V_2 S_2$

Where  $V_1$  = volume of insecticide required.

$S_1$  = strength of the commercial formulation

$V_2$  = volume of spray fluid required.

$S_2$  = strength of the spray fluid.

2. For granular application

$$\text{Quantity of chemical needed} = \frac{\text{Recommended a.i./ha}}{\% \text{ a.i in the formulation}} \times 100$$

### TECHNIQUES OF FUMIGATION FOR STORED GRAINS

**Precaution-** In the application of fumigants to grain streams, care should be taken that fumes are not inhaled. Liquid-type fumigants are especially hazardous because vapours may be given off before the grain enters the storage.

**Warning-** When grain fumigants are atomized or sprayed into closely confined spaces, or into a shallow space above the grain surface, the concentration of fumigant may exceed 2 percent by volume in air. Canister-type respirators will afford no protection under these conditions. It is better for the fumigator to remain outside and to apply the fumigant through an opening. If it is absolutely necessary for operators to enter such a space during fumigations, air-line or self-contained respirators should be worn.

**Dosage and Exposure:** Dosages of fumigants recommended for the mixing-in-grain. Dosage in fumigation of grain by direct mixing is modified by the kind of grain treated and the gas tightness of the structure. Wind forces, thermal expansion of the internal gas and changes in atmospheric pressure can also influence gas loss from storage structures

### **Methods of fumigation in godown**

**Direct mixing (vertical storage):** By this method, the fumigant is applied to the grain so that it is distributed as evenly as possible from the beginning of the treatment. Direct mixing is often employed when infestation is general throughout the mass and when there is access to the grain stream during filling or transfer from one bin to another. Only solid or liquid-type fumigants are used in this way. Aluminum phosphide tablets or pellets can be inserted in the grain stream by hand or with an automatic dispenser calibrated to deliver a dosage appropriate to the rate of loading in the bin and Calcium cyanide is usually discharged from an automatic applicator. Storage bins of the vertical type usually have manhole covers in the ceiling and these are usually closed immediately to prevent loss of fumigant

**Surface application (flat storage):** The surface application method has so far been used mainly with liquid type fumigants. The liquids are sprayed evenly over the top surface of the grain and the vapours slowly evolve and diffuse downward through the bulk. The carbon tetrachloride has given good distribution in grain in deep bins; carbon disulphide has been used in many countries.

**Large bulk Fumigation:** The liquid-type fumigants are usually applied to the surface of bulk grain by means of sprayers and the nozzles are removed to facilitate the rapid application of the liquid to the surface of the grain. A method for treatment of high vertical bins of grain by applying methyl bromide with carbon dioxide has been developed by Calderon and Carmi (1973). The carbon dioxide acts as a carrier and will take the methyl bromide down through the grain mass to the bottom of the bin. In flat storage units, in which the depth of the grain does not exceed 10 m (about 30 ft), tablets, pellets or sachets containing aluminium phosphide may be used. The fumigant is usually applied by probing into the grain

**Surface infestation:** With certain species of insects, such as the Indian meal moth, *Plodia interpunctella* infestation may be confined to the top of the grain. This problem cannot be solved by the usual method of surface application of fumigants because the vapours diffuse down through the grain. In silo bins or other storage units, which can be made air tight surface infestations can be treated with materials such as dichlorvos to obtain control. It should also be pointed out that incipient surface infestations of insects may be arrested by using pyrethrum, malathion or other approved materials applied as a fine mist in the space over grain. It should also be pointed out that incipient surface infestations of insects may be arrested by using pyrethrum, malathion or other approved materials applied as a fine mist in the space over grain. It should also be pointed out that incipient surface infestations of insects may be arrested by using pyrethrum, malathion or other approved materials applied as a fine mist in the space over grain.

**Hot spot fumigation:** Treatment of localised areas in a grain mass is often a useful technique for dealing with incipient infestation. These spots are usually recognised and defined by local rise in










temperature. Liquid type fumigants are applied through tubes. Aluminium phosphide tablets are the best materials in use. Enough fumigant is applied to maintain the required lethal concentration not only in the region of infestation, but also in the margins surrounding it for 1 or 2 m.

**Tent fumigation:** A tent (polythene or nylon impregnated with vinyl chloride) is constructed to cover sacks of grains. Tent can be conveniently stretched. The fumigant is introduced in to the tent through hoses connected to the preparing equipment. Most commonly HCN is used as fumigants

**Vaccum fumigation:** This is done in the case of cotton bales, imported products likely to be infested by insects, packaged food. The article to be fumigated is placed in a tight sealing steel chamber from which the air has been sucked out to produce a partial vacuum. A fumigant heated 120°F. Due to the partial vacuum the fumigant is able to penetrate the deeper layers of the bales. The reduction in the oxygen content due to partial vacuum forces the insects to breathe in toxic gas more readily. Today, the technique is used chiefly in plant quarantine work and for fumigating tobacco and other materials, such as compressed bales of jute bags and pressed dates which are difficult to penetrate at atmospheric pressure. Fumigants - Ethylene oxide/carbon dioxide mixture, Methyl bromide, Hydrogen cyanide.

**Fumigation for rodent control:** Most commonly employed chemical control measures include poison baits and fumigation. Initially poison baits are employed for control like Zinc phosphide (2%), Racumin bait, nor bromide etc. But to control the residual population of rodent's fumigation is necessary to kill more than 90% of population, otherwise they breed so fast that population reaches the same level within months. For this Fumigation with Aluminum phosphide tablets, 2 tablets of 0.6 g or half of 3g per burrow have been found effective. After introducing a tablet into live burrow, the opening is closed tight with soil. Soil moisture is essential to produce deadly phosphine gas.

**NON- INSECT PESTS ATTACKING IN FIELD/ GODOWN**

		
<b>Giant African Snail</b>	<b>Slug</b>	<b>Red Spider Mite</b>
		
<b>Root Knot Nematode</b>	<b>Millipede</b>	<b>House mouse</b>
		
<b>Black Drongo</b>	<b>House sparrow</b>	<b>House rat</b>

**DETERMINATION OF MOISTURE CONTENT OF GRAINS**



**Principle:** Removal of moisture from wet materials takes place by vaporization and it depends on the rate of heat and mass transfer which is related with two basic phenomenon namely vaporization of moisture from surface of material and movement of moisture from internal parts of materials to its surface. Movement of moisture takes place because of diffusion cell contraction and vapor pressure gradient.

**Requirements:** Electric balance, brown fuel moisture meter, stake moisture meter, indosaw universal moisture meter, oven, desiccators, moisture boxes

**Procedure:**

**Oven drying method**

1. Take the sample box and weigh with lid over it.
2. Put the sample in it (approximately 5-10 g)
3. Keep the sample in an oven at 105o C for 24 hrs
4. Take out the sample after 24 hrs and weigh it along with lid over it

Calculate the moisture content in percent with the following formula

**Brown dual distillation method**

Moisture content (wet basis) =  $\frac{\text{initial weight of sample} - \text{final weight of sample}}{\text{Initial weight of sample}}$

Moisture content (dry basis) =  $\frac{\text{initial weight of sample} - \text{final weight of sample}}{\text{Dry weight of sample}}$

**Procedure**

1. Arrange the instrument and settings
2. Take 100 g of material by weighing on the balance
3. Take 150 ml mineral oil (high bp) using jar
4. Take the grain and oil in a flask and keep it in the assembly
5. Supply the current and keep it for 30 minutes
6. Collect the condensed water in a graduated cylinder
7. Stop the supply when water collected in the cylinder is negligible
8. Take the cylinder and measure the reading which will give directly the wet basis moisture content

**Universal moisture content**

1. Arrange the instrument and set up
2. Take the sample and check the volume cup to be used from the mater
3. Fill the sample in the cup to top
4. Read out the pressure to be applied
5. Provide the compaction by means of a racket handle
6. Press the button provided at the top such that the countdown starts from 10
7. After the end of the countdown moisture content is displayed on the screen which gives the moisture content on weight basis
8. Repeat for three samples of same material

**Precautions**

1. The temperature and time for moisture removal are maintained properly
1. Clean and dry moisture boxes should be used for EXERCISEation
2. Condensed water should be collected properly and weighed

**METHODS OF GRAIN SAMPLING UNDER STORAGE CONDITIONS**

**Equipment Needed:** Deep bin compartment probe, Deepcup probe, Grain sieve with 1/12- or 3/16-inch round holes, Sample vials, Bin inspection forms, Temperature probe

**How to Sample:** While standing on the grain mass surface, push the probe into the grain mass at a slight angle. The top of the cup will open as the probe is pulled up and out of the grain, allowing grain to fill the cup. It is best to divide the grain surface into quarters and take at least three probes per quarter

section of grain mass. This will provide a good representative sample of the grain to allow inspection for the presence of insects, molds or excessively moist grain.

**Sampling Difficulties:** Overfilled grain bins are difficult to sample for insects or molds. Sometimes the only access points are through the bin wall, door or roof. Sample in the center of the grain mass as deeply as possible. Reach the bin wall if possible at two to three depths. Examine the Sample Place the grain sample in a specially designed weevil sieve (1½-inch diameter holes) if available and shake side to side at least 30 times to loosen any insects that may be in the grain. If a sieve is not available, place samples on a white piece of cloth for examination. Inspect the sample carefully for insects. It may be necessary to use a magnifying glass to see some of the smaller insects.

**Safety Precautions:** Bridged grain may result in a cave-in and subsequent suffocation of the workers. Bridged grain is caused when grain mats together, forming a false floor in the upper level of the grain mass. Persons falling through this bridged area are subject to suffocation.